

2100 New Bern Ave, Raleigh, NC 27610 P. 855-326-9112 / F. 855-326-9114

Date: Ne	eds by date:	Ship to: Patient Dilice 1st dose Dilice All doses								
Prescriber information										
Prescriber:		NPI:								
Supervising physician:		NPI:								
Address:										
Phone:		Fax:								
Office contact:										

Gastroenterolog		σv	Thoric.				T U.A.		
dastroenterology			Office contact:						
			F	Patient	informat	ion			
Name:			Sex:	□М	□F	DOB:	9	SSN:	
Address:			City:			State:	7	Zip:	
Phone:			Emerg	ency co	ntact:	Pł	none:		
Veight: Height:			Allergies:						
Primary language: ☐English ☐Spanish ☐Other:			Needs interpreter: ☐Yes ☐No						
	Please include	copies of	the patient	t's ins	urance	card or local	pharmacy inform	ation.	
					informa				
Diagnosis/ICD-10:	☐ K51. Ulcera	Icerative Colitis Other: ICD-10: ICD-10:							
	Test result: es □No		Test date:						
Prior failed thera	pies:	Duration				Duration			Duration
□Azathioprine			□Mesalamine				□Sulfasalazine		
□ Corticosteroids		□Methotrexate □					<u> </u>		
□Mercaptopurine			□Remicade □						
		Pre	scription infor	rmation	1			Quantity	Refills
⊒Cimzia®	200mgx2 Pre-filled syringe	\square Initial: Inject 400mg subcutaneously at weeks 0, 2, and 4					1 starter kit		
		☐Maintenance: Inject 400mg subcutaneously every 28 days					4 week supply		
□Dificid	200mg tablet	Take one tabl	Take one tablet by mouth twice daily						
Humira □ 40mg/0.8ml □ 40mg/0.4ml citrate	□Crohn's/UC Starter Kit	-	160mg subcutaneously day 1, then 80mg day 15					1 starter kit	
	□Pen	☐Maintenance: Inject 40mg subcutaneously every 14 days						4 week supply	
ree □Remicade	Vials □Initial:		: mg day 0, day 14, and day 42						
			Maintenance: Infuse mg every 8 weeks						
□Simponi	□100mg SmartJect □Initial:		ial: Inject 200mg subcutaneously week 0, then 100mg week 2					Loading dose	
	□100mg Pre-filled syringe							4 week supply	
¬c	□130mg vials	Initial: Infuse □260mg □390mg □520mg intravenously week 0							
□Stelara	□90mg Pre-filled syringe								
□Xeljanz	□10mg tablets	□ Initial: Take 10mg by mouth twice daily							
	☐5mg tablets	☐Maintenance: Take 5 mg by mouth twice daily							
-		☐ Maintenance: Take 10mg by mouth twice daily							
□Xifaxan	550mg tablets	Take one tablet by mouth three times daily					1 month supply		
□Other								1 month supply	

PHYSICIAN SIGNATURE REQUIRED

By signing below I authorize Josefs Pharmacy and its representatives to serve as a prior authorization designated agent in dealing with medical and prescription insurance companies.

Stamp signature not allowed, physician attests this is his/her legal signature.

Dispense as Written / Brand Medically Necessary

Substitution allowed

IMPORANT CONFIDENTIALITY STATEMENT: This message contains confidential, legally privileged, and/or proprietary information, intended only for the use of the named addressee above. If you are not the named addressee, you are hereby notified that any reading, retaining, disclosure, copying, distribution, or any action taken in reliance on the contents of these documents is strictly prohibited. If you have received this document in error, please notify Josefs Pharmacy immediately and destroy the transmitted information.